

# UC Davis

## UC Davis Previously Published Works

### Title

Draft Genome Sequences of Strains TAV3 and TAV4 (Verrucomicrobia: Opitutaceae), Isolated from a Wood-Feeding Termite, and In Silico Analysis of Their Polysaccharide-Degrading Enzymes.

### Permalink

<https://escholarship.org/uc/item/7220z8sc>

### Journal

Microbiology resource announcements, 9(2)

### ISSN

2576-098X

### Authors

Kotak, Malini  
Lin, Jonathan Y  
Isanapong, Jantiya  
et al.

### Publication Date

2020

### DOI

10.1128/mra.01192-19

Peer reviewed



# Draft Genome Sequences of Strains TAV3 and TAV4 (*Verrucomicrobia*: *Opitutaceae*), Isolated from a Wood-Feeding Termite, and *In Silico* Analysis of Their Polysaccharide-Degrading Enzymes

Malini Kotak,<sup>a</sup>  Jonathan Y. Lin,<sup>b</sup> Jantiya Isanapong,<sup>c</sup>  Jorge L. M. Rodrigues<sup>b,d</sup>

<sup>a</sup>Department of Biology, University of Texas, Arlington, Texas, USA

<sup>b</sup>Department of Land, Air and Water Resources, University of California, Davis, California, USA

<sup>c</sup>King Mongkut's University of Technology North Bangkok, Bangkok, Thailand

<sup>d</sup>Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA

**ABSTRACT** Here, we report the high-quality draft genome sequences of *Opitutaceae* sp. strains TAV3 and TAV4, which were isolated from the hindgut of the wood-feeding termite *Reticulitermes flavipes*. Using a combination of Illumina and PacBio sequencing, we constructed nearly complete assemblies totaling 5.84 and 5.91 Mbp in length for strains TAV3 and TAV4, respectively. In addition, we report an *in silico* analysis of potential lignocellulose-digesting enzymes present in these strains.

Wood-feeding termites are capable of efficiently digesting lignocellulose with the help of a wide variety of microorganisms residing in their hindgut (1, 2). Previously, five members of the bacterial family *Opitutaceae* (phylum *Verrucomicrobia*) were isolated from the hindgut of the wood-feeding termite *Reticulitermes flavipes* (3, 4). We used a combination of high-throughput sequencing to reconstruct the genome sequences of two isolates, strains TAV3 and TAV4, from the family *Opitutaceae* (subdivision 4).

The genomes of strains TAV3 and TAV4 were sequenced using Illumina MiSeq and PacBio platforms. For Illumina sequencing, single colonies of strains TAV3 and TAV4 were inoculated in liquid R2B medium from R2 agar plates inoculated from glycerol stocks kept in a  $-80^{\circ}\text{C}$  freezer. Liquid cultures were grown at  $25^{\circ}\text{C}$  for 1 week (EMD Chemicals, Billerica, MA), and genomic DNA was isolated using the MasterPure complete DNA and RNA purification kit (Epicentre Biotechnologies, Madison, WI) according to the manufacturer's instructions. Illumina TruSeq DNA libraries were prepared from paired-end 350-bp inserts and sequenced on the Illumina MiSeq platform with 75-bp paired-end reads. Illumina sequencing yielded 20,683,006 reads for strain TAV3 and 21,234,536 reads for strain TAV4, resulting in  $>270\times$  sequencing coverage for both strains.

For PacBio sequencing, the strains were cultured as mentioned previously, and the genomic DNA was extracted using the QIAamp DNA minikit (Qiagen, Valencia, CA) according to the manufacturer's instructions. Extracted DNA was quantified using the Qubit double-stranded DNA (dsDNA) high-sensitivity (HS) assay kit (ThermoFisher, San Jose, CA). The libraries were prepared using a SMRTbell template prep kit (Pacific Biosciences, Menlo Park, CA), and size selection for 20 kb was performed using the BluePippin system (Sage Scientific, Beverly, MA). The genomes of strains TAV3 and TAV4 were sequenced using the Sequel sequencing kit 2.1 and a 600-minute collection time. A total of 943,530 reads (subread  $N_{50}$ , 21,250 bp) were generated for strain TAV3, and a total of 1,093,102 reads (subread  $N_{50}$ , 18,750 bp) were generated for strain TAV4.

Reads were quality checked using FastQC (5), and a hybrid assembly was performed using reads from the Illumina and PacBio runs with Unicycler v0.4.6.0 (6). Default

**Citation** Kotak M, Lin JY, Isanapong J, Rodrigues JLM. 2020. Draft genome sequences of strains TAV3 and TAV4 (*Verrucomicrobia*: *Opitutaceae*), isolated from a wood-feeding termite, and *in silico* analysis of their polysaccharide-degrading enzymes. Microbiol Resour Announc 9:e01192-19. <https://doi.org/10.1128/MRA.01192-19>.

**Editor** David A. Baltus, University of Arizona

**Copyright** © 2020 Kotak et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jorge L. M. Rodrigues, [jmrodrigues@ucdavis.edu](mailto:jmrodrigues@ucdavis.edu).

**Received** 26 September 2019

**Accepted** 2 December 2019

**Published** 9 January 2020

parameters were used for all software unless otherwise specified. The draft genomes were annotated using the NCBI Prokaryotic Genome Annotation Pipeline and Prokka v1.12 (7, 8). The final assembly for strain TAV3 has a total length of 5,844,025 bp containing 4,976 genes predicted from 32 contigs, with an  $N_{50}$  value of 294,410 bp. Similarly, the TAV4 genome assembly contains 5,914,438 bp with 5,039 genes from 33 contigs, with an  $N_{50}$  value of 480,638 bp. Both genomes have a GC content of 60.9 mol% and 29 spanned (within scaffold) gaps which comprise 0.049% of the genome sequences.

A comprehensive *in silico* analysis of the carbohydrate-active enzyme (CAZy) profiles identified 433 (8.7% of total) and 431 (8.5% of total) genes in strains TAV3 and TAV4, respectively, as belonging to one of the CAZy families (9). The CAZy annotation identified 235 and 236 genes in TAV3 and TAV4, respectively, as glycoside hydrolases (GHs) belonging to 52 different families (10). TAV3 and TAV4 genomes also contain 83 and 85 glycosyltransferases (GTs), 8 and 7 polysaccharide lyases (PLs), 42 and 40 carbohydrate esterases (CEs), and 61 and 59 carbohydrate-binding modules (CBMs), respectively. Digestion of extracellular cellulose requires the action of secreted enzymes (11). A total of 104 CAZys in each of the strain TAV3 and TAV4 genomes contain signal peptides that suggest extracellular secretion of these proteins. These results show that strains TAV3 and TAV4 have the genomic potential to digest both cellulose and hemicellulose present in plant biomass.

**Data availability.** The whole-genome sequences and annotations for strains TAV3 and TAV4 have been deposited under the DDBJ/ENA/GenBank accession numbers [NZ\\_LXWT000000000](#) and [NZ\\_LXWU000000000](#). The NCBI BioProject and BioSample accession numbers for the TAV3 project are [PRJNA321366](#) and [SAMN04992812](#), respectively. The TAV3 raw reads were deposited in the SRA under accession numbers [SRR10174322](#) (PacBio) and [SRR3537542](#) (Illumina). The NCBI BioProject and BioSample accession numbers for the TAV4 project are [PRJNA321367](#) and [SAMN04992813](#), respectively. The TAV4 raw reads were deposited in the SRA under accession numbers [SRR10176387](#) (PacBio) and [SRR3537605](#) (Illumina).

## ACKNOWLEDGMENTS

High-throughput MiSeq sequencing services were performed at the University of Missouri DNA Core Facility. PacBio sequencing was performed at the DNA Technologies and Expression Analysis Core at the UC Davis Genome Center, supported by NIH shared instrumentation grant 1S10OD010786-01.

We thank Jeffery Demuth (University of Texas—Arlington) for encouraging discussions and Thomas Chrzanowski (University of Texas—Arlington) for letting us use his laboratory facility.

We declare no conflicts of interest.

## REFERENCES

- Brune A. 2014. Symbiotic digestion of lignocellulose in termite guts. *Nat Rev Microbiol* 12:168–180. <https://doi.org/10.1038/nrmicro3182>.
- Brune A, Dietrich C. 2015. The gut microbiota of termites: digesting the diversity in the light of ecology and evolution. *Annu Rev Microbiol* 69:145–166. <https://doi.org/10.1146/annurev-micro-092412-155715>.
- Stevenson BS, Eichorst SA, Wertz JT, Schmidt TM, Breznak JA. 2004. New strategies for cultivation and detection of previously uncultured microbes. *Appl Environ Microbiol* 70:4748–4755. <https://doi.org/10.1128/AEM.70.8.4748-4755.2004>.
- Isanapong J, Hambright WS, Willis AG, Boonmee A, Callister SJ, Burnum KE, Paša-Tolić L, Nicora CD, Wertz JT, Schmidt TM, Rodrigues JLM. 2013. Development of an ecophysiological model for *Diplosphaera colotermitum* TAV2, a termite hindgut *Verrucomicrobium*. *ISME J* 7:1803–1813. <https://doi.org/10.1038/ismej.2013.74>.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Cantarel BL, Coutinho PM, Rancurel C, Bernard T, Lombard V, Henrissat B. 2009. The Carbohydrate-Active EnZymes database (CAZy): an expert resource for glycogenomics. *Nucleic Acids Res* 37:233–238. <https://doi.org/10.1093/nar/gkn663>.
- Park BH, Karpins TV, Syed MH, Leuze MR, Uberbacher EC. 2010. CAZymes analysis toolkit (CAT): Web service for searching and analyzing carbohydrate-active enzymes in a newly sequenced organism using CAZy database. *Glycobiology* 20:1574–1584. <https://doi.org/10.1093/glycob/cwq106>.
- Blifernez-Klassen O, Klassen V, Doebe A, Kersting K, Grimm P, Wobbe L, Kruse O. 2012. Cellulose degradation and assimilation by the unicellular phototrophic eukaryote *Chlamydomonas reinhardtii*. *Nat Commun* 3:1214. <https://doi.org/10.1038/ncomms2210>.